

Quantum vacuum influence on pulsars spindown evolution

ARNAUD DUPAYS^{1,2,3,4(a)}, CARLO RIZZO⁵ and GIOVANNI FABRIZIO BIGNAMI⁶

¹ *Université de Toulouse, UPS, Laboratoire Collisions Agrégats Réactivité, IRSAMC - F-31062 Toulouse, France, EU*

² *CNRS, UMR 5589 - F-31062 Toulouse, France, EU*

³ *UPS-OMP, IRAP - Toulouse, France, EU*

⁴ *CNRS, UMR 5277 - F-31400 Toulouse, France, EU*

⁵ *Laboratoire National des Champs Magnétiques Intenses (UPR 3228, CNRS-INSA-UJF-UPS) F-31400 Toulouse Cedex, France, EU*

⁶ *IUSS, Istituto Universitario di Studi Superiori - Pavia, Italy, EU*

received 7 February 2012; accepted in final form 20 April 2012
published online 22 May 2012

PACS 97.60.Gb – Pulsars

PACS 12.20.Ds – Specific calculations

PACS 12.20.Fv – Experimental tests

Abstract – In this letter, we show that Quantum Vacuum Friction (QVF) resulting from the interaction between the magnetic dipole moment of a pulsar and its induced quantum vacuum magnetic dipole moment should play an important role in how a neutron star's spin period evolves. Taking into account this effect we show that magnetars could be understood as a natural evolution of standard pulsars. In particular, for the Crab pulsar, of which the true age is known, we present the first completely coherent time evolution for its period and braking index. For this pulsar and for the B1509-58 pulsar we also give the predicted values of the current first derivative of the braking index that we compare with existing experimental data. Our prediction provides a very important test to confirm or invalidate QVF.

 Copyright © EPLA, 2012

One of the most fundamental Quantum ElectroDynamics (QED) prediction concerns electric and magnetic properties of vacuum [1,2]. Following this idea, vacuum can be regarded as a standard medium with its own electromagnetic properties. Nevertheless, effects related to these properties are extremely weak and can only be tested in the presence of a very high electromagnetic field. For that reason, pulsars are very appropriate systems to look for evidences for vacuum properties [3,4]. Pulsars are fast-rotating neutrons stars, with a very high magnetic dipole moment tilted with respect to their rotational axis. Typically, their mass is of the order of the solar mass, and their radius of the order of 10 km. But the most interesting feature of these stars is certainly their magnetic field which can exceed 10^8 T on their surface [5]. Pulsars are thus suitable laboratories to study magnetic properties of the surrounding vacuum. In particular, a class of pulsars called magnetars [6] is characterized by a long spinning period and a magnetic field on the surface which is supposed to be even greater than the QED critical field 4×10^9 T [7,8]. At such a field non-linear properties of quantum vacuum

cannot be neglected. In general, the surface magnetic field of pulsars is inferred by comparing the measured period derivative with the one obtained using the energy loss rate given by the classical formula for a radiating rotating magnetic dipole. We believe that inferring a field exceeding the QED critical one using a pure classical formula (which neglects any QED effect) is questionable. Heyl *et al.* [9,10] have shown that the energy loss rate of a radiating rotating magnetic dipole, calculated by taking into account QED differs very little from the classical one. The use of a QED formula for the radiating dipole does not change in practice the inferred magnetic field on the surface of the pulsar.

Very recently, we have shown that a new energy loss process for pulsars should be taken into account, namely Quantum Vacuum Friction (QVF) resulting from the interaction between the non-stationary magnetic dipole moment of the star and its induced quantum vacuum magnetic dipole moment [11]. In the case of magnetars, with very high magnetic fields, we have shown that the QVF is the dominating energy loss process and has important consequences. In particular, the inferred value of the magnetic field no long exceeds the QED critical field for all known magnetars.

^(a)E-mail: arnaud.dupays@irsamc.ups-tlse.fr




Fig. 1: P - \dot{P} diagram for the currently known pulsars. Pulsars located in the upper right area (gray) have an energy loss process dominated by QVF. The Crab pulsar is marked as a circle filled with a dot.

In this letter we consider some astrophysical consequences of QVF. We show that QVF plays a crucial role in neutron star spindown evolution. Taking into account this effect, we show that magnetars can simply be understood as a natural evolution state of typical pulsars, similar to the Crab pulsar. For the Crab pulsar, of which the true age is known, we present the first completely coherent time evolution for its period and braking index. Let us first summarize the main results concerning QVF in highly magnetized neutron stars. Considering a neutron star rotating in vacuum, we denote by R its radius and B_0 the magnitude of the magnetic field at its surface. If P is the spinning period of the neutron star and θ the inclination angle of its magnetic dipole moment with respect to the rotation axis of the star, one can show that for $B_0 \leq 10^{10}$ T, the QVF energy loss rate \dot{E}_{qv} is given by [11]

$$\dot{E}_{qv} \simeq \alpha \left(\frac{3\pi^2}{4} \right) \frac{\sin^2 \theta}{B_c^2 \mu_0 c} \frac{B_0^4 R^4}{P^2}, \quad (1)$$

where $B_c = m_e^2 c^3 / e \hbar \simeq 4.4 \times 10^9$ T is the QED critical field, α the fine structure constant, μ_0 the vacuum permeability and c the light velocity *in vacuo*. Total energy loss rate of the star \dot{E} can thus be obtained by adding the classical dipolar radiative energy loss rate \dot{E}_r given by

$$\dot{E}_r = \left(\frac{128\pi^5}{3} \right) \frac{\sin^2 \theta}{\mu_0 c^3} \frac{B_0^2 R^6}{P^4}. \quad (2)$$

It is important to stress that the spinning period dependence is not the same for the classical radiative process and for the QVF. The ratio of classical to QVF losses decreases like $1/P^2$ for large P , so that QVF becomes more important for slowly rotating neutron stars. Let us now study how QVF drastically changes the evolution of a pulsar. The evolution of a pulsar in period P and slowdown \dot{P} is usually described in a logarithmic P - \dot{P} diagram as shown in fig. 1, where we have plotted all of the currently

known pulsars (~ 1700) for which P and \dot{P} have been measured [12]. On this diagram, we have also indicated lines of constant magnetic field in the case where only the classical radiative energy loss process is considered (dotted lines) together with the case for which QVF is taken into account (solid curves). These lines are obtained by matching the loss in rotational energy \dot{E} with $\dot{E}_r + \dot{E}_{qv}$. Since \dot{E} is given by

$$\dot{E} = 4\pi^2 I \dot{P} P^{-3}, \quad (3)$$

where a neutron star moment of inertia of $I = 10^{45}$ g cm² is assumed, we get

$$\dot{P} = \left(\frac{32\pi^3}{3I} \right) \frac{B_0^2 R^6}{\mu_0 c^3 P} + \alpha \left(\frac{3}{16I} \right) \frac{B_0^4 R^4 P}{B_c^2 \mu_0 c}, \quad (4)$$

where $\sin \theta$ is taken, as usual, equal to 1. The first term in eq. (4) corresponds to the classical contribution, which gives the asymptotic behaviour for small P . The second term is the correction due to QVF, which becomes important for large P . In fig. 1 we clearly distinguish two different evolution areas: pulsars located in the upper right area of the figure (gray) have an energy loss process dominated by QVF, whereas pulsars in the lower left area have classical magnetic dipole emission as the main energy loss. Defining the borderline between both areas as

$$\frac{d \log \dot{P}}{d \log P} = 0, \quad (5)$$

we get

$$\frac{\ddot{P} P}{\dot{P}^3} = 0 \quad (6)$$

Since $P \neq 0$, this equation implies that $\ddot{P} = 0$ and thus, the borderline between both areas corresponds to pulsars having a constant slowdown \dot{P} . Assuming that the pulsar's magnetic field to remain constant during its evolution, we can describe the evolution of the star following the line on which it is situated today. Let us first ignore QVF. Since the rotational period of a pulsar increases with time, the natural evolution of these stars corresponds to a fall towards the lower right area of the figure along a given straight dotted line, depending of the value of their magnetic field. In this case, pulsars located in the upper right area can exhibit surface magnetic fields above the quantum critical field. Since these stars cannot be connected by a straight line of constant magnetic field to other pulsars with lower rotational period, we have to assume that these stars are very young, even if their rotational period is already high. For that reason, this class of pulsars, called magnetars, seems to be very different in nature from other standard pulsars. To understand the structure of these stars and the physical process involved which should give rise to such a huge surface magnetic field remains an open question. If we take into account QVF, we see that all magnetars are now connected by a modified line of constant magnetic field to pulsars with lower rotational period. Remarkably enough, their surface

magnetic field becomes lower than the critical quantum field. We can then conclude, as far as QVF exists, that magnetars could be understood as a natural evolution of standard pulsars. Such a very important astrophysical consequence of quantum vacuum effect shows that testing the validity of the QVF theory would be crucial in the future to understand pulsars physics. It is also important to stress that with magnetic fields below the critical quantum field, the current explanation for the emission properties of Anomalous X-ray Pulsars (AXPs) and Soft Gamma-ray Repeaters (SGRs) cannot be applied to our model [13,14], and so, this emission remains an open question.

Another interesting result of taking QVF into account in magnetized rotating neutron stars is a completely coherent time evolution of pulsars, as we will show in the following. Let us illustrate this point in detail with the Crab pulsar (the one marked as a circle filled with a dot in fig. 1). Situated in the center of the Crab Nebula which is the remnant of the Super Novae SN1054, the Crab pulsar is certainly one of the most well-known pulsars [15]. A very interesting feature of this pulsar is that historical records indicate its true age to be ~ 955 yr. This indication, together with accurate measurements of its current spin period P , first and second time derivative of the spin period, respectively P_1 and P_2 , drastically constrains its rotational evolution dynamics. Rewriting eq. (4) as

$$\dot{P} = \frac{K_r}{P} + K_{qv}P, \quad (7)$$

where $K_r = \dot{E}_r \times (P^4/4\pi^2 I)$ and $K_{qv} = \dot{E}_{qv} \times (P^2/4\pi^2 I)$, the age of a pulsar τ is given by

$$\tau = \int_{t_0}^t dt' = \int_{P_i}^P \frac{dP'}{\frac{K_r}{P'} + K_{qv}P'}, \quad (8)$$

where t denotes the current date and P_i the initial spin period at $t = t_0$. We obtain

$$\tau = \frac{1}{2K_{qv}} \ln \left(\frac{K_r + K_{qv}P^2}{K_r + K_{qv}P_i^2} \right). \quad (9)$$

If the initial value of the period P_i would be known together with the today value P , eq. (9) would give the age of a pulsar taking into account QVF and classical dipole radiation emission.

If $K_{qv} \ll K_r$ eq. (9) becomes

$$\tau \simeq \frac{1}{2K_r} (P^2 - P_i^2). \quad (10)$$

From eq. (7) we have in this case $K_r \simeq \dot{P}P$, so that

$$\tau \simeq \frac{P}{2\dot{P}} \left(1 - \left(\frac{P_i}{P} \right)^2 \right). \quad (11)$$

Finally, making the assumption that $P_i \ll P$ eq. (11) becomes

$$\tau \simeq \frac{P}{2\dot{P}}, \quad (12)$$




Fig. 2: Spin period of the Crab pulsar as a function of time when QVF is taken into account (full line) and when QVF is neglected (dotted line). $t=0$ corresponds to current time (*i.e.* “today”).

which is the expression usually employed to estimate the characteristic age of a pulsar taking into account only the classical dipole radiation emission.

In general, eq. (9) can be written as

$$\tau = \frac{1}{2K_{qv}} \ln \left(\frac{1 + \frac{K_r}{K_{qv}P^2}}{\frac{K_r}{K_{qv}P_i^2} + \frac{P_i^2}{P^2}} \right). \quad (13)$$

Even in the case that $P_i \ll P$ eq. (13) cannot be reduced to an equation like the classical eq. (12) since $P_i \ll P$ does not always mean that $\frac{P_i^2}{P^2} \ll \frac{K_r}{K_{qv}P^2}$.

Once the age of a pulsar is known together with P and \dot{P} , our eq. (9) can be used to infer the P_i value exactly as the classical eq. (10) using the value of the magnetic field B_0 obtained inverting eq. (4) to calculate K_r and K_{qv} . In particular this also applies to the case of AXPs/SGRs magnetars. The ages of the supernova remnants independently give a measure of their ages, which suggest that they are young. The P_i value inferred from the experimental measured age taking into account QVF and classical dipole radiation emission is generally different from the one inferred using eq. (13) which only takes into account classical dipole radiation emission.

To illustrate this last point let study the case of the Crab pulsar. Inverting eq. (9) (respectively, eq. (11)) we obtain the spin period as a function of time when classical and QVF energy loss processes are taken into account (respectively, when QVF is neglected). Both cases are presented in fig. 2. In this figure, $t=0$ corresponds to current time (*i.e.* “today”). As already discussed, since the age of this pulsar is known, we can estimate the initial spin period P_i .

Using the classical expression given by eq. (10) we obtain $P_i = 16$ ms, whereas we get $P_i = 18$ ms when QVF is taken into account. Since the energy loss process of the Crab pulsar is dominated by classical radiation (see fig. 1), we are not surprised that the global time evolution

of this star does not drastically change when QVF effects are included. Nevertheless, the most important difference between both calculations is that the one taking into account QVF is the only one which provides a spin period evolution completely coherent with current observations, in particular with the current value of the braking index, as we will discuss in the following.

The braking index, n , is a fundamental dimensionless quantity given by [16]

$$n = 2 - \frac{P\ddot{P}}{\dot{P}^2}, \quad (14)$$

where P is the spin period of the pulsar and \dot{P} (respectively, \ddot{P}), denotes the first (respectively, the second) derivative of P with respect to time. This parameter describes the rate at which a pulsar loses rotational energy. From this expression we see that the borderline between gray and white plotted in fig. 1 corresponds to pulsars having a braking index equal to 2 ($\ddot{P} = 0$). Pulsars located in the upper right area of the figure (gray) have a braking index lower than 2, whereas braking index of pulsars in the lower left area is higher. In the following we study the time evolution of n for $t > 0$. We denote by P_0 the current spin period, and by P_1 and P_2 , the current values of the first and the second derivative of P with respect to time. The current value of n is

$$n_0 = 2 - \frac{P_0 P_2}{P_1^2}. \quad (15)$$

At this stage it is important to stress that the constant value $n = 3$ is expected for pure classical radiation models, whereas the current measured value is for example, $n_0 = 2.51$ for the Crab pulsar [15]. Very recently, Espinoza *et al.* have even found $n_0 \simeq 1$ for magnetar J1734-3333 [17]. This is exactly what we have predicted for a magnetar [11], however Espinoza *et al.* suggest an alternative interpretation based on an hypothetical grows of the magnetic field. In any case, these discrepancies show that pure classical radiation models would not lead to a realistic picture for pulsar's time evolution. Adding QVF effects, solution of eq. (7) becomes, for $t \geq 0$

$$p^2(t) = \left(\frac{K_r}{K_{qv}} + P_0^2 \right) e^{2K_{qv}t} - \frac{K_r}{K_{qv}}. \quad (16)$$

This expression allows us to calculate the first and the second derivative of P with respect to time, and to match $\dot{P}(0)$ with P_1 and $\ddot{P}(0)$ with P_2 , obtaining

$$K_{qv} = \frac{P_0 P_2 + P_1^2}{2P_0 P_1} \quad (17)$$

and

$$K_r = \frac{P_0 P_1^2 - P_2 P_0^2}{2P_1}. \quad (18)$$

Finally, the time evolution of the spin period is given by

$$P(t) = \sqrt{\frac{2P_0 P_1}{A} (e^{At} - 1) + P_0^2} \quad (19)$$




Fig. 3: Braking index of the Crab pulsar as a function of time when QVF is taken into account (full line) and when QVF is neglected (dotted line). $t = 0$ corresponds to current time.

with

$$A = \left(\frac{P_2}{P_1} + \frac{P_1}{P_0} \right). \quad (20)$$

Using eq. (19), we can calculate the braking index

$$\begin{aligned} n(t) &= 1 - \left(\frac{P_0 P_2}{P_1^2} - 1 \right) e^{-At} \\ &= 1 - (1 - n_0) e^{-At}. \end{aligned} \quad (21)$$

The braking index time evolution for the Crab pulsar is presented in fig. 3. When QVF is taken into account, the braking index is a decreasing function with respect to time which is consistent with a global rotation slowdown.

From eq. (21) we can predict the current value of the first derivative of n given by

$$\dot{n}(0) = A(1 - n_0). \quad (22)$$

As far as we know, there are only 8 stars in the ATNF catalogue [12], our reference for pulsar data, for which P_0 , P_1 , P_2 have been measured giving $1 < n_0 < 3$. For two of them the P_3 , the third derivative of P with respect to time, have been measured. These stars are the Crab pulsar [15] and the B1509-58 pulsar [18]. For these two stars the value of $\dot{n}^c(0)$ can be calculated from the first derivative of eq. (14) leading to

$$\dot{n}^c(0) = (4 - n_0) \frac{P_2}{P_1} - \frac{P_0 P_3}{P_1^2}. \quad (23)$$

This value can be compared with our estimation given by eq. (22). For the Crab pulsar we get $\dot{n}(0) \sim -1.8421 \times 10^{-11} \text{ s}^{-1}$, a value 100 times away from the observation since $\dot{n}^c(0) \sim 1.5676 \times 10^{-13} \text{ s}^{-1}$. However we would like to point out that taking QVF into account allows to perform an exact calculation even for a non-constant n , something that the classical approximation does not allow. For the B1509-58 pulsar we get

$\dot{n}(0) \sim -1.3550 \times 10^{-11} \text{ s}^{-1}$, a value in perfect agreement with the calculated one $\dot{n}^c(0) \sim -1.2500 \times 10^{-11} \text{ s}^{-1}$. These results show that new measurements of $\dot{n}(0)$ are needed to improve statistics and thus provide a very important test to confirm our or invalidate QVF predictions. Of course we are fully aware of the difficulties of this long-term pulsar measurement as outlined in [19]. We still trust that pulsars will allow QVF measurement to be performed and thus give further support to QED.

In this paper we have shown that QVF should play an important role in neutron star spindown evolution. Taking into account this effect, we have shown that the surface magnetic field of magnetars becomes lower than the critical quantum field, so that magnetars could be simply understood as a natural evolution of standard pulsars. For pulsars for which the braking index is known, QVF provides a completely coherent spindown history, as illustrated with the Crab pulsar. For this pulsar and for the B1509-58 pulsar we have also given the predicted value of the current first derivative of the braking index, suggesting a very important test to confirm or invalidate QVF. Both issues are very important. Observations of QVF in pulsars would confirm the existence of one of the very few macroscopic effects of quantum vacuum, while if QVF is invalidated this would indicate for the first time that the usual QED model of the quantum vacuum used for calculating the QVF effect needs to be mended.

REFERENCES

- [1] EULER H. and KOCKEL B., *Naturwissenschaften*, **23** (1935) 246; HEISENBERG W. and EULER H., *Z. Phys.*, **98** (1936) 714.
- [2] RIKKEN G. L. J. A. and RIZZO C., *Phys. Rev. A*, **63** (2001) 012107.
- [3] DUPAYS A., ROBILLIARD C., RIZZO C. and BIGNAMI G. F., *Phys. Rev. Lett.*, **94** (2005) 161101.
- [4] DUPAYS A., RIZZO C., RONCADELLI M. and BIGNAMI G. F., *Phys. Rev. Lett.*, **95** (2005) 211302.
- [5] LYNE A. and GRAHAM-SMITH F., *Pulsar Astronomy* (Cambridge University Press) 1989.
- [6] KOUVELIOTOU C., DUNCAN R. C. and THOMPSON C., *Sci. Am. Int. Ed.*, **288** (2003) 24.
- [7] IBRAHIM A. I., SWANK J. H. and PARKE W., *Astrophys. J. Lett.*, **584** (2002) L17.
- [8] BIGNAMI G. F., CARAVEO P. A., DE LUCA A. and MEREGHETTI S., *Nature*, **423** (2003) 725.
- [9] HEYL J. S. and HERNQUIST L., *Phys. Rev. D*, **55** (1997) 2449.
- [10] HEYL J. S. and HERNQUIST L., *J. Phys. A: Math. Gen.*, **30** (1997) 6485.
- [11] DUPAYS A., RIZZO C., BAKALOV D. and BIGNAMI G. F., *EPL*, **82** (2008) 69002.
- [12] MANCHESTER R. N., HOBBS G. B., TEOH A. and HOBBS M., *Astron. J.*, **129** (2005) 1993 (astro-ph/0412641) (see also the web address <http://www.atnf.csiro.au/research/pulsar/psrcat>).
- [13] QIAO G. J., XU R. X. and DU Y. J., (2010) arXiv:1005.3911v1 [astro-ph.HE].
- [14] NG C.-Y. and KASPI V. M., (2010) arXiv:1010.4592v1 [astro-ph.HE].
- [15] LYNE A. G., PRITCHARD R. S. and SMITH F. G., *Mon. Not. R. Astron. Soc.*, **265** (1993) 1003.
- [16] MANCHESTER R. N. and TAYLOR J. H., *Pulsars* (Freeman, San Francisco) 1977.
- [17] ESPINOZA C. M., LYNE A. G., KRAMER M., MANCHESTER R. N. and KASPI V. M., *Astrophys. J. Lett.*, **741** (2011) L13.
- [18] KASPI V. M., MANCHESTER R. N., SIEGMAN B., JOHNSTON S. and LYNE A. G., *Astrophys. J.*, **422** (1994) L83.
- [19] HOBBS X., LYNE A. G. and KRAMER M., *Mon. Not. R. Astron. Soc.*, **402** (2010) 1027.